

Heights of Models of ZFC and the Existence of End Elementary Extensions

Andrés Villaveces

February 1, 2008

Abstract

The existence of End Elementary Extensions of models M of ZFC is related to the ordinal height of M , according to classical results due to Keisler, Morley and Silver. In this paper, we further investigate the connection between the height of M and the existence of End Elementary Extensions of M . In particular, we prove that the theory ‘ $ZFC + GCH +$ there exist measurable cardinals $+$ all inaccessible non weakly compact cardinals are possible heights of models with no End Elementary Extensions’ is consistent relative to the theory ‘ $ZFC + GCH +$ there exist measurable cardinals $+$ the weakly compact cardinals are cofinal in ON ’. We also provide a simpler coding that destroys GCH but otherwise yields the same result.

I wish to thank my advisor, Kenneth Kunen, for many helpful conversations and comments, always full of interesting insights. I also wish to thank Ali Enayat for very helpful discussions about some of the topics treated in this article, Sy Friedman for interesting questions and comments related to the class forcing construction around Theorem 3.1, Mirna Džamonja and Arnie Miller for various helpful discussions.

1 Introduction.

Let $(M, E), (N, F)$, etc. denote models of ‘enough set theory.’ The central notion of extension we use in this article is the well known ‘end elementary

extension'. A model (N, F) **end extends** (M, E) iff for every $a \in M$, the sets $a_E = \{b \in M \mid bEa\}$ and $a_F = \{b \in N \mid bFa\}$ are the same. In other words, elements of M are not enlarged by the extension from M to N .

Let $(\mathcal{E}_M, \prec_e)$ denote the structure of all non-trivial elementary end extensions ('ees') of M (a model of set theory), together with the relation $\prec_e$ (we write ' $\mathcal{A} \prec_e \mathcal{B}$ ' if and only if $\mathcal{B}$ is an elementary *end* extension of $\mathcal{A}$). $\prec_e$ is an ordering on $\mathcal{E}_M$. The kind of ordering relation that $\prec_e$ is on $\mathcal{E}_M$ depends heavily on certain structural features of M . Notice that structures like $(\mathcal{E}_M, \prec_e)$ need not be well-founded: as Kaufmann notes in [Ka 83], if κ is weakly compact, $(\mathcal{E}_{\mathcal{R}(\kappa)}, \prec_e)$ has infinite descending chains.

In this paper, we concentrate on problems of **existence** of end elementary extensions of specific models of ZFC, and we study the relationship between the height of those models and the possibility to obstruct the existence of ees. In a forthcoming work, we will concentrate on the study of chains in $(\mathcal{E}_M, \prec_e)$, and their connection with large cardinal properties of M .

We consider the following general question:

Question 1 *How does the structure of M affect the structure of $(\mathcal{E}_M, \prec_e)$?*

The earliest results toward a solution to Question 1 were obtained by Keisler, Silver and Morley in [KeSi 70] and [KeMo 68]. All their theorems addressed the specific question of the **existence** of ees (when is $\mathcal{E}_M \neq 0$?). They provided answers for the two following cases.

Theorem 1.1 *(Keisler, Morley [KeMo 68]) Let M be a model of ZFC, $\text{cof}(M) = \omega$. Then $\mathcal{E}_M \neq 0$. (This takes care in particular of all countable models of ZFC.)*

Theorem 1.2 *(Keisler, Silver [KeSi 70]) If M is of the form $\mathcal{R}(\kappa)$, where κ is a weakly compact cardinal, then for all $S \subset M$, $\mathcal{E}_{(\mathcal{R}(\kappa), \in, S)} \neq 0$.*

In light of those early results, it is natural to ask to what degree does the **height** of M determine the structure of $\mathcal{E}_M$. We begin by the basic question: does the height of M determine the **existence** of an eee?

Here are the two main results of this paper.

Theorem 2.2: The theory ‘ZFC + $\exists \lambda$ measurable + $\forall \kappa(\kappa$ inaccessible not weakly compact $\rightarrow \exists$ transitive $M_\kappa \models ZFC$ such that $o(M) = \kappa$ and $\mathcal{E}_M^{wf} = 0$)’ is consistent relative to the theory ‘ZFC + $\exists \lambda$ measurable’.

Theorem 3.1: The theory ‘ZFC + GCH + $\exists \lambda(\lambda$ measurable) + $\forall \kappa[\kappa$ inaccessible not weakly compact $\rightarrow \exists$ transitive $M_\kappa \models ZFC$ such that $o(M) = \kappa$ and $\mathcal{E}_M^{wf} = 0]$ ’ is consistent relative to the theory ‘ZFC + $\exists \lambda(\lambda$ measurable) + the weakly compact cardinals are cofinal in ON’.

The ‘conclusion’ in both theorems is that the existence of transitive models of the form $\mathcal{R}(\kappa)$ which have eees, yet the inner model $M_\kappa \subset \mathcal{R}(\kappa)$ does not have any well-founded eees (that is, the ‘global’ negative answer to the Height Problem, later abbreviated as the ‘NED’ property) is consistent with fairly reasonable large cardinal axioms.

The proofs in both cases use codings of the obstructions to the existence of eees **by an appropriate inner model**. Although the results look similar, the codings used in both cases differ strongly: in the first theorem, the places where GCH holds or fails provide the main tool for the coding; in the second case, since one needs to obtain models where GCH holds everywhere, one cannot use anymore such a device for the coding. In that case, forcing nonreflecting stationary sets at the appropriate levels to a model previously freed of any of those sets does the trick.

The notation we use is standard. We restrict ourselves to the case of *transitive* M . Following two different traditions, we freely switch between the two notations ‘ V_κ ’ and ‘ $\mathcal{R}(\kappa)$ ’ when we denote the set of objects of the universe

of rank less than κ . Given a model M , $o(M)$ denotes the ordinal height of M .

2 The Height Problem.

When M is definable, the problem is trivial, by the following

Proposition 2.1 *If M is definable in $\mathcal{R}(\kappa)$, and $\mathcal{R}(\kappa)$ has an eee, then M has an eee.*

Proof: Let N elementarily end extend $\mathcal{R}(\kappa)$. Since $o(M) = \kappa$, M^N , the interpretation of the definition of M in N , properly (end) extends M . It is easy to verify that we actually have $M \prec_e M^N$. $\square$

Of course, this is far from settling the general problem

Question 2 (Height Problem) *Do $\mathcal{E}_{\mathcal{R}(\kappa)} \neq 0$ and $o(M) = \kappa$ imply together that $\mathcal{E}_M \neq 0$?*




Figure 1: The Height Problem

As Theorem 2.2 shows, it is quite possible that this question have a negative answer: there are models of set theory where a certain $\mathcal{R}(\kappa)$ has ees, yet some transitive submodel with the same height has no ees. This can even

be obtained in a quite homogeneous way: the ‘extendability property’ only holds at weakly compact cardinals.

Still, if we add new axioms to the theory, the answer to the question may become positive. This is the case, for example, when $V = L$ holds in M , the question becomes trivial, since in that case the only transitive submodel of height κ of an $\mathcal{R}(\kappa)$ is $\mathcal{R}(\kappa)$ itself.

Therefore, the answer to the Height Problem is independent of ZFC.

2.1 Consistency of a global negative answer.

We will next present some of the possible situations for answers to a version of the problem. Let $G(\lambda)$ denote any large cardinal property preserved under Easton-type extensions where the iteration process is not carried too often (measurability, etc.). More precisely, we mean here properties unaffected by Easton iterations which only act at certain **successor** cardinals.

Definition 1 *Let κ be an inaccessible cardinal. We say that κ is NON-END-DETERMINING (or for short $NED(\kappa)$) if and only if there is a transitive $M_\kappa \models ZFC$ such that $o(M) = \kappa$ and $\mathcal{E}_M^{wf} = 0$.*

Thus, if $NED(\kappa)$, there may be models of the form $\mathcal{R}_\kappa$ which have ees, yet certain inner models of them do not have well-founded ees. Observe that the NED property depends strongly on the universe where κ is being considered.

Theorem 2.2 *The theory ‘ $ZFC + \exists \lambda G(\lambda) + \forall \kappa (\kappa \text{ inaccessible not weakly compact} \rightarrow NED(\kappa))$ ’ is consistent relative to the theory ‘ $ZFC + \exists \lambda G(\lambda)$ ’.*

We will devote the remainder of this section to proving Theorem 2.2 and to draw some corollaries from it. The proof of Theorem 2.2 is based on a forcing construction which produces codes for various sets which witness the non existence of ees at the desired cardinals. The ‘ $\exists \lambda G(\lambda)$ ’ clause is

added to the theory to point out that the forcing can be done in such a way that large cardinal properties $G(\lambda)$ (at least, those not destroyed by Easton iterations which only act at certain **successor** cardinals) are consistent with the negative answer to Question 2.

In the next lemma, we introduce the forcing construction, and we show that both weak compactness and non weak compactness are preserved in the extension.

Lemma 2.3 *Let M be a model of $ZFC+GCH$, and let $M[G]$ be a forwards Easton extension obtained by adding α^{++} subsets to every successor inaccessible α . Then M and $M[G]$ have the same weakly compact cardinals.*

Proof: Let first κ be weakly compact. Notice that κ is not in the domain of the Easton iteration: it only acts at successor inaccessibles. Let ϕ be Σ_1^1 (say $\phi \equiv \exists X \psi(X)$, ψ first order). Let

$$1 \Vdash_{\mathbb{P}} (\forall \alpha < \kappa \exists X \subset \alpha [(\alpha, <, \tau) \models \psi(X)]),$$

for some $\mathbb{P}$ -name τ in V such that $1 \Vdash \tau \subset \kappa$. Then, by the Π_1^1 -indescribability of κ in M , there are N , $\mathbb{P}^*$, τ^* such that

$$(\mathcal{R}(\kappa), \in, \tau, \mathbb{P}) \prec_e (N, \in, \tau^*, \mathbb{P}^*).$$

Now, in M , $\mathbb{P}^*$ is an Easton forcing which (since $\mathbb{P}^*$ does not act on κ) can be seen as a product

$$\mathbb{P}^* = \mathbb{P} \times \mathbb{P}'.$$

A usual argument shows that $\mathbb{P}'$ is $(< \kappa^+)$ -closed. Then, $\mathbb{P}^*$ does not add any new subsets to κ . Now, in M , we have

$$1 \Vdash_{\mathbb{P}^*} \exists X \subset \kappa ((\kappa, <, \tau, X) \models \psi).$$

So, since $M^{\mathbb{P}^*} = (M^{\mathbb{P}'})^{\mathbb{P}}$,

$$1 \Vdash_{\mathbb{P}'} \exists \text{ some } \mathbb{P}\text{-name } \sigma [1 \Vdash_{\mathbb{P}} ((\kappa, <, \tau, \sigma) \models \psi)].$$

Hence, by the κ -closure of $\mathbb{P}'$, a fusion argument in the style of Silver provides a $\mathbb{P}$ -name σ in M such that

$$1 \Vdash_{\mathbb{P}} ((\kappa, <, \tau, \sigma) \models \psi).$$

So, κ is also weakly compact in $M[G]$.

Now, if κ is *not* weakly compact in the universe, there is a κ -Aronszajn tree T . There are two cases:

- (i) κ is Mahlo: then $\mathbb{P}$ is the product of a κ -Knaster and a κ^+ -closed forcing; thus

$$1 \Vdash T \text{ is } \kappa\text{-Aronszajn,}$$

whereby κ cannot be weakly compact in $M[G]$.

- (ii) κ is not Mahlo. Then, there is a club C of singular cardinals in κ , $C \in M$. None of these singular cardinals becomes regular in $M[G]$; clearly C is also a club in κ in $M[G]$. So, κ remains non-Mahlo in the extension. $\square_{2.3}$

Having shown this preservation property of $\mathbb{P}$, we are ready to complete the

Proof of Theorem 2.2: Assume that GCH and $\exists \lambda(G(\lambda))$ both hold in V . Let $V[G]$ be a forward Easton extension like in the previous Lemma. Then, V and $V[G]$ have the same weakly compact cardinals. If κ is *not* weakly compact (in V or $V[G]$), then we need to find a transitive M_κ in $V[G]$ with no eee and $o(M_\kappa) = \kappa$. There are three cases:

- 1) κ is a successor inaccessible: then let $M_\kappa = \mathcal{R}(\kappa)$. This M_κ cannot be end elementary extended: when we have $N \succ_e M_\kappa$, κ turns out to be inaccessible in N . But this implies that the inaccessible are unbounded in κ , and this is incompatible with the fact that κ is a successor inaccessible.

- 2) κ is a non-Mahlo limit inaccessible: then fix C a club in κ with no regular cardinals, and let M_κ code C by the powers of the successor inaccessibles below κ . This M_κ is obtained as follows: write κ as an increasing sequence $(\lambda_\alpha)_{\alpha < \kappa}$, where the α 's run over the successor inaccessibles below κ . Now, set $M_\kappa =$ the extension of $\mathcal{R}(\kappa)$ where α^{++} subsets of α are added only when $\lambda_\alpha \in C$. Clearly, we have $\mathcal{R}(\kappa)^V \subset M_\kappa \subset \mathcal{R}(\kappa)^{V[G]}$ and $o(M_\kappa) = \kappa$. Since M_κ codes C , it cannot have any eee (if it had one, say N , then κ would be a singular cardinal in $N!$).
- 3) κ is Mahlo: then let T be a κ -Aronszajn tree in V . Since T remains κ -Aronszajn in $V[G]$, it is enough to let M_κ encode T similarly to part 2; this new M_κ cannot be elementarily end extended, for it would then provide a κ -branch to T in $V[G]$, contradicting the fact that T is Aronszajn. This completes our proof. $\square_{2.2}$

Corollary 2.4 *The Height Problem may have a negative answer.*

Proof: In the model $V[G]$ constructed in Theorem 2.2, let κ be the first inaccessible such that $\mathcal{E}_{\mathcal{R}(\kappa)} \neq 0$. Results in [KeSi 70] show that κ is not weakly compact. Then, by Theorem 2.2, there exists a transitive model M_κ of ZFC which is of height κ and has no well-founded ees. This provides a negative answer to the Height Problem. $\square$

It is interesting to observe the following, in connection to Theorem 2.2.

Proposition 2.5 *If κ is weakly compact, then for all transitive $M \subset \mathcal{R}(\kappa)$ of height κ , $\mathcal{E}_M \neq 0$.*

Proof: It suffices to observe that if κ is weakly compact, then $(\mathcal{R}(\kappa), \in, S)$ has ees, for all $S \subset \mathcal{R}(\kappa)$. Take any (transitive) $M \subset \mathcal{R}(\kappa)$ of height κ . Then $(\mathcal{R}(\kappa), \in, M)$ has a well founded eee $(N, \in, M')$. Clearly,

- i $M \subseteq_e M'$: by endness, all the ‘new’ elements go ‘on top’ of M , and

- ii $M \prec M'$: given any sentence ϕ , $\mathcal{R}(\kappa) \models (M \models \phi)$ if and only if $N \models (M' \models \phi)$. This implies $M \prec M'$ by the transitivity of N . $\square$

The ‘opposite’ problem is still open, for cardinals κ which are inaccessible not weakly compact:

Open Question 1 *Is the theory ‘ZFC + large cardinals + $\forall \kappa$ inaccessible not weakly compact $((M \models \text{ZFC} \wedge o(M) = \kappa \wedge \mathcal{E}_{\mathcal{R}(\kappa)} \neq 0) \rightarrow \mathcal{E}_M \neq 0)$ ’ consistent?*

In other words, is it consistent to have large cardinals and simultaneously a *globally* positive answer to the Height Problem? Up to now, the only way to get globally positive answers to the Height Problem is with $V = L$ or similar axioms.

Remark: An easy modification of the construction given for Theorem 2.2 provides models for the Height Problem only at certain cardinals.

We basically have two extreme situations here: the Height Problem under the very restrictive $V = L$ (trivial positive answer since there are no strictly inner models), and the Height Problem in the presence of Large Cardinals (negative answers). It is natural to ask

Question 3 *What happens ‘in between’ $V = L$ and Large Cardinal Axioms? More specifically, what is the situation for models of the form $L[0^\#]$ or $L[\mu]$ (where μ is a measure in the usual sense)?*

The bottom line here is that in the presence of large cardinals, there are enough sharps which allow encoding the parameters on a model of the form $\mathcal{R}(\kappa)$, for κ non weakly compact. So, a situation similar to that of Theorem 2.2 would be obtained via a κ -closed poset: in $L[\mu]$, if κ is not weakly compact, then there is a model M of height κ with no ees. Situations similar to these are the subject of the final section.

3 A new coding: keeping GCH true throughout the forcing.


In the previous section, the construction of models for $NED(\kappa)$ used Mc Aloon's coding method: coding (in inner models of the generic extension) the Aronszajn trees or the clubs of singulars needed to destroy ees by using the successor inaccessibles where GCH holds/fails as the coding device. Naturally, if one wants to obtain a similar result while *keeping* GCH true, one must code the construction in a completely different way.

Theorem 3.1 *The theory ‘ $ZFC + GCH + \exists\lambda(\lambda \text{ measurable}) + \forall\kappa[\kappa \text{ inaccessible not weakly compact} \rightarrow NED(\kappa)]$ ’ is consistent relative to the theory ‘ $ZFC + \exists\lambda(\lambda \text{ measurable}) + \text{the weakly compact cardinals are cofinal in } ON$ ’.*

This answers a question that Sy Friedman asked me during the Tenth Latin American Mathematical Logic Symposium in Bogotá. The proof of this theorem consists of a two step forcing, followed by a construction of models of height κ with no ees in the same way as in the end of the proof of Theorem 2.2: coding in M_κ an object that makes it impossible for M_κ to have ees. (Depending on the case, a κ -Aronszajn tree or a club of singulars.)

The two forcing constructions provide enough coding tools to complete the proof in a way analogous to how coding was used to prove Theorem 2.2. The idea is to use the existence/non-existence of nonreflecting stationary sets on α^{++} (those which will be called $E_{\alpha^{++}}^\alpha$ -sets in the construction), for α successor inaccessible, instead of the continuum function (cardinals where GCH holds/fails), to do the coding.

We start with a model of ‘ $ZFC + GCH + \exists\lambda(\lambda \text{ measurable}) + \text{the weakly compacts are cofinal in } ON$ ’. We first force the non-existence of ‘ E -sets’, and then force back the existence of them on successor inaccessibles. We will then be able to construe the inner models we need inside the final generic model by putting in it **exactly** the E -sets one needs to carry the coding.

Figure 2: Coding M by adding only **some** E -sets (the $\square$ s)

Definition 2 (E_κ^θ , nonreflecting stationary sets) Let θ be a regular cardinal, and $\kappa > \theta$, with $\text{cof}(\kappa) > \aleph_1$. We mean by E_κ^θ that

$$\exists A \subset \{\delta < \kappa \mid \text{cof}(\delta) = \theta\}, \text{ } A \text{ stationary in } \kappa \text{ such that}$$

$$\forall \text{ limit } \alpha < \kappa (A \cap \alpha \text{ is not stationary in } \alpha).$$

Abusing the notation a bit, we sometimes say that the set A is an ' E_κ^θ -set', or even an ' E -set'.

Proof of Theorem 3.1: Start with M , a model of 'ZFC + GCH + $\exists \lambda(\lambda \text{ measurable}) + \forall \alpha \exists \mu > \alpha(\mu \text{ weakly compact})$ '.

Let $\mathbb{P}$ be the Easton product of the Lévy collapses $\mathbb{L}_i = \text{Coll}(\kappa_i^+, < \theta_i)$ of cardinals $< \theta_i$ to κ_i^+ , where $i \in \text{ORD}^M$, and the sequence $\langle \langle \kappa_i, \theta_i \rangle \mid i \in \text{ORD}^M \rangle$ is defined as follows: κ_0 is the first successor inaccessible cardinal, and θ_0 is the first weakly compact cardinal; if κ_i, θ_i are defined, then κ_{i+1} is the first successor inaccessible *after* θ_i and θ_{i+1} is the first weakly compact cardinal after κ_{i+1} . For limit i , κ_i is the first successor inaccessible after $\sup\{\kappa_j \mid j < i\}$, and θ_i is the first weakly compact cardinal after κ_i .

More precisely, $p \in \mathbb{P}$ iff p is a function such that

- i $p(i) \in \mathbb{L}_i$,
- ii for all $i \in \text{dom}(p)$, for all σ inaccessible such that $i < \sigma$ implies $\theta_i < \sigma$, $\text{dom}(p) \cap \sigma$ is bounded in σ .

$\mathbb{P}$ is ordered in the usual way.

For every κ , $\mathbb{P}$ factors into three forcings: $\mathbb{P} \approx \mathbb{P}_{<\kappa} * \mathbb{P}_\kappa * \mathbb{P}_{>\kappa}$, where $\mathbb{P}_{<\kappa}$, $\mathbb{P}_\kappa$, $\mathbb{P}_{>\kappa}$ mean respectively the portions of the iteration below κ , at κ , and above κ . By Easton's Lemma and *GCH* in M , $\mathbb{P}$ collapses the cardinals in $[\kappa_i^+, \theta_i)$ to κ_i^+ (thus, in the generic extension, θ_i becomes κ_i^{++}), and preserves all the other cardinals. In particular, all the κ_i s remain successor inaccessible in the extension, as the reader can easily check.

Claim: For all i , for all stationary $S \subset \{\beta < \kappa_i^{++} \mid \text{cof}(\beta) = \kappa_i\}$ in $V^\mathbb{P}$, there exists σ such that $\text{cof}(\sigma) = \kappa_i^+$ and $S \cap \sigma$ is stationary. (We say that the stationary reflection property holds for κ_i and κ_i^{++} at κ_i^+ .)

To show this, it is enough to look at $V^{(\mathbb{P}_{\leq \theta_i})}$, since this coincides with $V^\mathbb{P}$ up to the level κ_{i+1} . $V^{(\mathbb{P}_{\leq \theta_i})}$ can be seen as $(V^{\mathbb{L}_i})^{\mathbb{P}_{<\theta_i}}$. Since the stationary reflection property holds in V for κ_i and θ_i^+ at κ_i^+ (by Baumgartner [Ba 76]), we have that the stationary reflection property for κ_i and $\theta_i = \kappa_i^{++}$ (at κ_i^+ necessarily) also holds in $V^{\mathbb{L}_i}$. On the other hand, $|\mathbb{P}_{<\theta_i}| < \kappa_i^+ < \theta_i$; hence if S is a stationary subset of $\{\beta < \kappa_i^{++} \mid \text{cof}(\beta) = \kappa_i\}$, and $S \in (V^{\mathbb{L}_i})^{\mathbb{P}_{<\theta_i}}$, then there is $T \subset S$ stationary, $T \in V^{\mathbb{L}_i}$. By the stationary reflection property in $V^{\mathbb{L}_i}$, there exists some $\gamma < \theta_i$, with $\text{cof}(\gamma) = \kappa_i^+$, such that $T \cap \gamma$ is stationary in $V^{\mathbb{L}_i}$. $T \cap \gamma$ is still stationary in $(V^{\mathbb{L}_i})^{\mathbb{P}_{<\theta_i}}$, since $|\mathbb{P}_{<\theta_i}| < \kappa_i^+$. (Since whenever $p \Vdash \dot{\tau}$ club in γ ($\text{cof } \gamma = \kappa_i^+$), we have that $\dot{\tau}$ already is a club in γ in $V^{\mathbb{L}_i}$.) All this ends the proof of the Claim.

So, the stationary reflection property holds in $V^\mathbb{P}$ for all κ_i , θ_i . This ends the first stage of the proof. We have obtained a model $V_1 = V[G]$, for G $\mathbb{P}$ -generic, where $E_{\theta_i}^{\kappa_i}$ -sets do NOT exist, for any $i \in \text{ORD}$.

For the second step of the proof, we will simplify matters by writing ' V ' instead of ' V_1 ' and calling our forcing $\mathbb{P}$ again. We need to force $E_{\kappa^{++}}^\kappa$ -sets

into κ^{++} , for each successor inaccessible κ . We will achieve this goal by ‘piecewise adding’ the E -sets.

The proof that such E -sets can be added is an Easton iteration. We present here a variant of the construction by Cummings, Džamonja and Shelah in [CuDžSh 95] of a generic extension of a model of ‘ $GCH + \exists \lambda (\lambda \text{ measurable})$ ’, where a regular cardinal θ ‘strongly non-reflects’ at λ_i , where $i < \theta$, $\text{cof}(\lambda_i) = \lambda_i < \lambda_i^+ < \theta$. Their construction yields in particular $E_\theta^{\lambda_i}$ -sets in the generic extension (see Lemma 2 and Theorem 1 of [CuDžSh 95]).

The Building Blocks: given any successor inaccessible cardinal κ , let $\mathbb{P}_\kappa$ consist of conditions $p = (\alpha_p, E_p, C_p)$, where $\alpha_p < \kappa^{++}$, $\text{cof}(\alpha_p) = \kappa^+$, $E_p \subset \alpha_p$,

$$C_p : S_{\kappa^+}^{\alpha_p} \rightarrow \bigcup_{\beta < \alpha_p} \mathcal{P}(\beta)$$

where $S_{\kappa^+}^{\alpha_p} = \{\beta < \alpha_p \mid \text{cof}(\beta) = \kappa^+\}$, and for all $\beta \in S_{\kappa^+}^{\alpha_p}$, $C_p(\beta)$ is a club on β . We also require that for all $\beta \in S_{\kappa^+}^{\alpha_p}$, $C_p(\beta) \cap E_p = \emptyset$, and $E_p \subset S_{\kappa^+}^{\alpha_p}$. So, E_p behaves like an ‘approximation’ to a non-reflecting stationary subset of $S_{\kappa^+}^{\kappa^{++}}$. We define the ordering of $\mathbb{P}_\kappa$ by

$$q \leq p \iff \begin{cases} \text{(a)} & \alpha_q \geq \alpha_p, \\ \text{(b)} & E_q \cap \alpha_p = E_p, \\ \text{(c)} & C_q \upharpoonright S_{\alpha_p}^{\kappa^+} = C_p. \end{cases}$$

We need only worry at ordinals of cofinality κ^+ :

Lemma 3.2 *If $\text{cof}(\alpha) \in [\aleph_1, \kappa^+)$, then there is a club $C \subset \alpha$ such that for all $\gamma \in C$, $\text{cof}(\gamma) < \kappa$.*

Proof: Fix α with $\aleph_1 \leq \text{cof}(\alpha) \leq \kappa$, and let $f : \text{cof}(\alpha) \rightarrow \alpha$ be increasing and continuous. Then all the γ ’s in the domain of f are $< \kappa$. As f is continuous increasing, $\text{ran}(f)$ is a club in α , and for every $\beta \in \text{ran}(f)$, we have $\text{cof}(\beta) = \text{cof}(f^{-1}(\beta)) < \kappa$. $\square$

Let now G be $\mathbb{P}_\kappa$ -generic, and let $E_G = \bigcup_{p \in G} E_p$.

Claim: E_G is a nonreflecting stationary subset of $S_{\kappa^{++}}^\kappa$.

To show that E_G is nonreflecting, let $\alpha < \kappa^{++}$ be of cofinality κ^+ . Then $\alpha < \alpha_p$ for some $p \in G$, and thus $C_{\alpha_p}(\alpha) \cap E_{\alpha_p} = 0$. But by (b) in the definition of $\leq$, $E_G \cap \alpha_p = E_{\alpha_p}$, $C_{\alpha_p}(\alpha) \subset \alpha < \alpha_p$. So, $C_{\alpha_p}(\alpha)$, which is a club on α , does not intersect E_G : $C_{\alpha_p}(\alpha) \cap E_G = (C_{\alpha_p}(\alpha) \cap \alpha_p) \cap E_G = C_{\alpha_p}(\alpha) \cap (\alpha_p \cap E_G) = C_{\alpha_p}(\alpha) \cap E_{\alpha_p} = 0$.

To see that E_G is stationary, let Γ be a canonical name for E_G , and suppose that $p \Vdash (\tau \text{ club of } \kappa^{++} \wedge \tau \cap \Gamma = 0)$. Get then a decreasing κ -chain of conditions $p > p_0 > p_1 > \dots > p_\xi > \dots$, $p_\xi \in \mathbb{P}_\kappa$, such that for $\xi < \kappa$, $p_\xi \Vdash \delta_\xi \in \tau$ and $\alpha_{p_\xi} < \delta_\xi < \alpha_{p_{\xi+1}}$. Let $q = (\sup_{\xi < \kappa} \alpha_{p_\xi}, \bigcup_{\xi < \kappa} E_{p_\xi}, \bigcup_{\xi < \kappa} C_{p_\xi})$. This q is not a condition, since the cofinality of its support is $< \kappa^+$. But we just have to take any condition r that extends q in a natural way.

[How? Just let $r = (\sup_{\xi < \kappa} \alpha_{p_\xi} + \kappa^+, \bigcup_{\xi < \kappa} E_{p_\xi} \cup \{\alpha^*\}, \bigcup_{\xi < \kappa} C_{p_\xi} \cup \hat{C})$, where

- i $\alpha^* = \sup_{\xi < \kappa} \alpha_{p_\xi}$, and
- ii $\hat{C}(\beta)$ is any club on β with $\min(\hat{C}(\beta)) > \alpha^*$, for $\beta \in S_{\kappa^+}^{\alpha^* + \kappa^+} \setminus \alpha^*$.

It is not difficult to check that this $r \in \mathbb{P}_\kappa$.]

Then, the limit of the δ_ξ 's ($= \alpha^*$) is forced by r to belong to the club τ . On the other hand, $\alpha^* \in E_r$ and $r \Vdash \alpha^* \in \Gamma$. But this contradicts the fact that $p \Vdash (\tau \text{ club} \wedge \tau \cap \Gamma = 0)$.

Claim: $\mathbb{P}_\kappa$ is $(< \kappa^+)$ -closed.

This Claim is established by the same argument as the previous one, although taking now sequences of arbitrary lengths below κ^+ .

Remark: Clearly, $\mathbb{P}_\kappa$ is not $(< \kappa^{++})$ -closed, since $\mathbb{P}_\kappa$ adds κ^{++} -sequences.

The $\mathbb{P}_\kappa$'s correspond thus to the 'Building Blocks' of our construction. The remaining part consists of the iteration through all the successor inaccessible cardinals, and of the proof that measurables are preserved.

Claim: $|\mathbb{P}_\kappa| \leq \kappa^{++}$.

(Just observe that in V_1 , $\kappa^{++} \cdot 2^{\kappa^+} \cdot 2^{\kappa^+} = \kappa^{++}$.)

3.1 The Iteration.

The iteration is in the style of Backward Easton. The supports are bounded below regular cardinals. We add new subsets only at successor inaccessible stages. Formally, this corresponds to defining $\mathbb{P}_{\leq \alpha}$ as the forcing up to stage α and $\dot{Q}_\alpha \in V^{\mathbb{P}_\alpha}$ as the forcing at stage α . Set $\dot{Q}_\alpha = \{0\}$, if α is not a successor inaccessible, and $\dot{Q}_\alpha = (\mathbb{P}_\alpha)^{V_{<\alpha}}$, otherwise, where $V_{<\alpha}$ stands for $V^{\mathbb{P}_{<\alpha}}$. By the κ^+ -closure of the forcing $\mathbb{P}_\kappa$, this iteration yields a model of *ZFC*. In order to finish the proof, we need only show that the measurability of λ is preserved by this iteration.

Let λ be a measurable cardinal in V , and let $j : V \rightarrow M$ be the ultrapower map arising from a normal measure U on κ . We first observe that it is enough to prove that λ remains measurable in the extension by $\mathbb{P}_{<\lambda}$, the rest of the forcing being $(< \lambda^+)$ -closed, by the previous claim. (The power set of λ is not changed by $\mathbb{P}_{\geq \lambda} \approx \mathbb{P}_{> \lambda}$.)

As is well-known, j has the following properties:

1. $\text{crit}(j) = \lambda$.
2. ${}^\lambda M \subset M$.
3. $\lambda^+ < j(\lambda) < j(\lambda^+) < \lambda^{++}$.
4. $M = \{j(F)(\lambda) \mid F \in V \wedge \text{dom}(F) = \lambda\}$.

We want to prove that j ‘lifts’ to an embedding $\tilde{j} : V[G] \rightarrow N \subset V[G]$ (which we shall also denote by ‘ j ’, abusing notation), where G is $\mathbb{P}$ -generic over V , thereby automatically ensuring that λ is also measurable in $V[G]$. The idea is to prolong the generic G to a generic H for the forcing $j(\mathbb{P}_{<\lambda})$.

We start by comparing $\mathbb{P}_{<\lambda}$ to $j(\mathbb{P}_{<\lambda})$. $j(\mathbb{P}_{<\lambda})$ is an iteration defined in M , forcing the existence of non-reflecting stationary sets on each $S_\kappa^{\kappa^{++}}$, for each κ successor inaccessible $< j(\lambda)$. If we compute the iteration $j(\mathbb{P}_{<\lambda})$ up to stage λ , we get exactly $\mathbb{P}_{<\lambda}$. We can now compute a generic extension $M[G]$ of M using the V -generic filter, since $\mathbb{P}_{<\lambda}$ -generics over V are also generic over M . Since $|\mathbb{P}_{<\lambda}| < \lambda$, every canonical $\mathbb{P}_{\leq\lambda}$ -name for a λ -sequence of ordinals is in M , so in $V[G]$. Thus, we have

Claim: $V[G] \models {}^\lambda(M[G]) \subset M[G]$.

In $M[G]$, we will prolong G to a $j(\mathbb{P}_{<\lambda})$ -generic. Call the ‘remainder forcing’ $\mathbb{R} = \mathbb{R}_{\lambda, j(\lambda)}$, that is, $j(\mathbb{P}_{<\lambda}) = \mathbb{P}_{<\lambda} * \mathbb{R}$. $\mathbb{R}$ is $(< \lambda_M^+)$ -closed in $M[G]$; hence, by the previous claim, $\mathbb{R}$ is also $(< \lambda_M^+)$ -closed in $V[G]$. In $M[G]$, the forcing $\mathbb{R}$ is $j(\lambda)$ -cc and has size $j(\lambda)$. There are then at most $j(\lambda)$ maximal antichains in $\mathbb{R}$ in the model $M[G]$, as $[j(\lambda)^{j(\lambda)} = j(\lambda)]^M$, by elementarity and since $|\mathbb{P}_{<\lambda}| < \lambda$. In $V[G]$, we can enumerate those antichains as $\langle A_\alpha : \alpha < \lambda^+ \rangle$, since $\lambda^+ < j(\lambda) < j(\lambda^+) < \lambda^{++}$. Using the closure to meet all these antichains, it is clear that in $V[G]$, we can build H which is $\mathbb{R}$ -generic over $M[G]$. Letting $G^+ = G * H$, G^+ is $j(\mathbb{P}_\lambda)$ -generic over M . We define $j : V[G] \rightarrow M[G^+]$ by $j(\dot{\tau}^G) = j(\dot{\tau})^{G^+}$. This is a well-defined elementary embedding, as follows from the following fact, whose proof uses the Truth Lemma and the elementarity of j .

Fact: Let $k : M \rightarrow N$ be an elementary embedding between two transitive models of ZFC. Let $\mathbb{P} \in M$ be a forcing notion, let $k(\mathbb{P}) = \mathbb{Q}$, and suppose that G is $\mathbb{P}$ -generic over M and H is $\mathbb{Q}$ -generic over N . Let also $k''G \subset H$. Then the definition $k(\dot{\tau}^G) = k(\dot{\tau})^H$ for every $\dot{\tau} \in M^\mathbb{P}$ gives a well-defined elementary embedding $k : M[G] \rightarrow N[H]$, which extends $k : M \rightarrow N$ and is such that $k(G) = H$.

This ends the proof of measurability of λ in $V[G]$, and thus the proof of Theorem 3.1: At this point, to witness the non-end-determining property of the appropriate κ ’s, exactly the same argument used in the proof of Theorem 2.2 works. Instead of using the failure/non-failure of GCH at the successor

inaccessibles below κ to code the corresponding object (club of singulars or κ -Aronszajn tree), here, the coding is achieved by the inclusion/non-inclusion of the corresponding E -set (see figure 3). $\square_{3.1}$

4 Ees of Inner Models.

In this final section, we push the study of the connections between the existence of ees and the structure of the basic model one step further: We look at inner models of some specific models related to $0^\sharp$ or to measures, and we look at some instances of the ‘wider model problem’.

In the last section, we remarked that for $L[\mu]$, the situation is similar to the one obtained via Theorem 2.2: if κ is not weakly compact, then there is a model M of height κ with no ees. Contrasting this, we have the following fact for the inner model L^M , in presence of $0^\sharp$.

Proposition 4.1 *If $M \models ‘0^\sharp \text{ exists}’$, and M is a set model of ZFC, then $\mathcal{E}_{L^M} \neq 0$.*

Proof: The existence of $0^\sharp$ in M allows to code ees of L^M . $\square$

The converse to this is clearly false: just take $M = L_\kappa$ such that (for example), κ is weakly compact. Then, $\mathcal{E}_{L^M} \neq 0$, yet $M \models ‘0^\sharp \text{ does not exist}’$.

We have the following situation, asking the same question for *well founded* ees (compare the following result and Proposition 4.1, and to Kunen’s result in Kaufmann [Ka 83]).

Proposition 4.2 *Suppose that M satisfies*

- i *M is countable, and*
- ii *$M \models ‘0^\sharp \text{ exists}’$,*

and that M has minimal ordinal height among all the models which have simultaneously the properties **i** and **ii**. Then, $\mathcal{E}_{L^M}^{wf} = 0$.

Proof: We can use a Σ_2^1 formula to say that the height of a certain model M is α and that M thinks that $0^\sharp$ exists: let $\psi(\alpha) \equiv \text{'}\exists \text{ transitive } M(o(M) = \alpha \wedge M \models 0^\sharp \text{ exists})\text{'}$. This is clearly Σ_2^1 . Now, we also have by Löwenheim-Skolem that $ZFC \vdash [\exists \alpha \psi(\alpha) \rightarrow \exists \alpha < \omega_1 \psi(\alpha)]$. The formula $\psi(\alpha)$ relativises down to models N of ZFC when α^N is countable. So, working in V , we can fix M , a transitive model such that $M \models \text{'}0^\sharp \text{ exists'}$, $o(M) = \delta$, and $\delta < \omega_1$ is the least possible among the heights of such models. By Keisler-Morley, we know that $\mathcal{E}_{L^M} \neq 0$, and thus

$$L^M = L_\delta \models ZFC + \text{'inaccessibles are cofinal in } ORD\text{'}$$

Suppose then that $\mathcal{E}_{L^M}^{wf} \neq 0$; let $\gamma < \omega_1$ be such that $L_\gamma \succ L_\delta$. We then have that $L_\gamma \models \aleph_\delta = \delta$. A collapse of δ to ω does the trick: working in L_γ , let $\mathbb{P} = \text{Coll}(\omega, \delta)$, and let G be $\mathbb{P}$ -generic over L_γ . So, on one hand, in $L_\gamma[G]$, δ is countable, and on the other hand, $\psi(\delta)$ holds in V . Then, $\psi(\delta)$ also holds in $L_\gamma[G]$, by Σ_2^1 -absoluteness. But then, since L_γ and $L_\gamma[G]$ have the same ordinals, $\psi(\delta)$ holds in L_γ . Finally, by elementarity, $L_\delta \models \exists \theta(\psi(\theta))$. This contradicts the minimality of δ . $\square$

So far, the Height Problem has only been looked at for *inner* models of a given $\mathcal{R}(\kappa)$ which is known to have non trivial ees. Yet a natural question arises concerning ‘wider’ models. More specifically, we have the following

Question 4 (ees for wider models) *Let $\mathfrak{A}$ be an inner model, $\mathcal{E}_{\mathcal{R}(\kappa)^\mathfrak{A}} \neq 0$, $M \supset \mathcal{R}(\kappa)^\mathfrak{A}$, $M \models ZFC$, $o(M) = \kappa$, $|\kappa| > \omega$. When is $\mathcal{E}_M \neq 0$?*

In such generality, there is no simple answer to this question. Yet, we have that in particular, when M is a generic extension of $\mathcal{R}(\kappa)^\mathfrak{A}$, obtained via a set-forcing $\mathbb{P}$, M must also have non trivial ees. This is easily established




Figure 3: The ‘wider models problem’

by observing that if $N \in \mathcal{E}_{\mathcal{R}(\kappa)^{\mathfrak{A}}}$, and $M = \mathcal{R}(\kappa)^{\mathfrak{A}}[G]$, then $N[G] \succ_e M$. This was implicitly used in the proof of Theorem 2.2. There, we did not really have a set-forcing. Yet, the decomposition properties of Easton forcing acting only on successor inaccessible cardinals imply that in that case the forcing extension necessarily has eees.

An example of a CLASS forcing that destroys the property of $\mathcal{R}(\kappa)$ having eees can be obtained from Boos’s Easton style forcing construction from [Bo 74]. This was pointed out to me by Ali Enayat. Boos’s forcing, a variation on a thin-set forcing notion by Jensen, allows one to efface the Mahlo property:

Theorem 4.3 (Boos - [Bo 74]) *If M is a transitive model of $ZFC + GCH$, and κ is κ -Mahlo in M , then for each $\alpha < \kappa$ there is a forcing notion $\mathbb{B}_\alpha$ such that $M^{\mathbb{B}_\alpha} \models \text{‘}\kappa \text{ is strongly } \alpha\text{-Mahlo but not } (\alpha + 1)\text{-Mahlo’}$.*

Using the $\mathbb{B}_1$ from this theorem, any weakly compact cardinal in M becomes the first Mahlo in a generic extension. So, we have

Corollary 4.4 *There are Easton-type forcings which do not preserve the property of having eees: let κ be a weakly compact cardinal. Then, although we know by Keisler and Silver [KeSi 70] that $\mathcal{E}_{\mathcal{R}(\kappa)^M} \neq 0$, we have $\mathcal{E}_{\mathcal{R}(\kappa)^M[G]} = 0$, for G $\mathbb{B}_1$ -generic over $\mathcal{R}(\kappa)^M$.*

Remarks: 1) Boos's construction —a variant of a construction by Jensen— is an Easton-type iteration. By [KeMo 68], if κ is the first Mahlo cardinal, $\mathcal{R}(\kappa)$ cannot have any ees. The weakly compact κ of the corollary (in M) becomes the first Mahlo in the extension by $\mathbb{B}_1$.

2) Of course, if $\mathbb{B}_n$ is used instead of $\mathbb{B}_1$ here, then κ becomes the first n -Mahlo cardinal. The corresponding $\mathcal{R}(\kappa)^M[G]$ cannot either have elementary end extensions: if $N \succ_e \mathcal{R}(\kappa)^M[G]$, then $N \models \forall \lambda \exists C \subset \lambda (C \text{ club and } C \text{ does not contain any } (n-1)\text{-Mahlos})$. But then, this is also true about κ , and κ would not be n -Mahlo.

References.

- (Ba 76) BAUMGARTNER, J. *A new class of order types*, Annals of Mathematical Logic 9 (1976), 187-222.
- (BeJeWe 83) BELLER, A., JENSEN, WELCH *Coding the Universe*, Cambridge Lecture Series, 1983.
- (Bo 74) BOOS, W. *Boolean Extensions which efface the Mahlo property*, The Journal of Symbolic Logic, vol. 39, no 2, June 74, pages 254-268.
- (CuDžSh 95) CUMMINGS, J., DŽAMONJA, M., SHELAH, S. *A consistency result on weak reflection*, accepted by Fund. Math.
- (En 84) ENAYAT, A. *On certain Elementary Extensions of Models of Set Theory*, Trans. Amer. Math. Soc., 1984.
- (En ∞) ENAYAT, A. *Counting Countable Models of Set Theory*, preprint.
- (Ka 83) KAUFMANN, M. *Blunt and Topless Extensions of Models of Set Theory*, J. Symb. Logic, 48, 1983, 1053-1073.
- (KeMo 68) KEISLER, H.J., MORLEY, M. *Elementary Extensions of Models of Set Theory*. Israel J. of Math., vol. 6, 1968.

(KeSi 70) KEISLER, H.J., SILVER, J. *End Extensions of Models of Set Theory*, Proc. Symp. Pure Math. 13 1970 (177-187).